

OUTP-97-63P

27th November 1997

hep-th/9712058

The Spectral Dimension of Non-generic Branched Polymer Ensembles

JOÃO D. CORREIA¹ AND JOHN F. WHEATER²

*Department of Physics, University of Oxford
Theoretical Physics,
1 Keble Road,
Oxford OX1 3NP, UK*

Abstract. We show that the spectral dimension on non-generic branched polymer models with susceptibility exponent $\gamma > 0$ is given by $d_s = 2/(1 + \gamma)$. For those models with $\gamma < 0$ we find that $d_s = 2$.

PACS: 04.60.Nc, 5.20.-y, 5.60.+w

Keywords: conformal matter, quantum gravity, branched polymer, spectral dimension

¹e-mail: j.correia1@physics.ox.ac.uk²e-mail: j.wheater1@physics.ox.ac.uk

The spectral dimension d_s is an important measure of the dimensionality of the manifold ensembles appearing in quantum gravity. In particular much effort has recently been put into its determination for discretized two-dimensional quantum gravity ensembles. The spectral dimension is defined by a random walk which leaves a fixed vertex at $t = 0$ and at every step is allowed to move from its present position to one of its neighbouring vertices with uniform probability. After t steps the probability that the walk has returned to the initial point is given by

$$P(t) \approx \frac{\text{const}}{t^{d_s/2}} \quad (1)$$

provided that $t \gg 1$ (to negate discretization effects) and that $t \ll N^\Delta$ where N is the number of vertices in the graph and Δ some exponent (to avoid finite size effects). The spectral dimension has been investigated numerically for many different values of the central charge c [1] and it has been calculated analytically for the generic branched polymer model [2] and found to be $d_s = \frac{4}{3}$ in excellent agreement with the numerical results at large c . We refer the reader to refs [1] and [2] for detailed discussion of the definition of d_s and the ensemble averages involved. In this letter we extend the results of [2] to the non-generic branched polymers.

We start by recalling the essential features of the branched polymer models [3]. A general branched polymer (BP) model has grand canonical partition function $\mathcal{Z}(z)$ satisfying

$$\mathcal{Z}(z) = z(1 + \sum_{n=1}^{\infty} \alpha_n \mathcal{Z}(z)^n) \quad (2)$$

where the α_n are constants. Iterating this equation shows that $\mathcal{Z}(z)$ is the generating function for the set of all rooted trees $\mathcal{B}$ made up of links and vertices of all coordination numbers $n + 1$ such that $\alpha_n \neq 0$,

$$\mathcal{Z}(z) = \sum_{A \in \mathcal{B}} z^{N_A} w_A. \quad (3)$$

The number of links in A is denoted by N_A and the weight w_A of A is given by

$$w_A = \prod_{v \in A} \alpha_{n(v)-1} \quad (4)$$

where v runs over all vertices of A and $n(v)$ denotes the coordination number of v . We denote the set of all polymers whose first vertex has coordination number $n + 1$ by $\mathcal{B}^n$; clearly $\mathcal{B} = \mathcal{B}^0 \cup \mathcal{B}^1 \cup \mathcal{B}^2 \cup \dots$ ($\mathcal{B}^0$ contains the polymer consisting of a single link). Any graph $A \in \mathcal{B}^n$ has a set of constituents $A_1, \dots, A_n$ obtained by severing the links connecting the first vertex to the rest of the graph (see fig.1). Note that




Figure 1: The constituents $A_1, \dots, A_n$ of a branched polymer $A \in \mathcal{B}^n$.

for $A \in \mathcal{B}^n$

$$w_A = \alpha_n \prod_{i=1}^n w_{A_i} \quad (5)$$

and

$$N_A = 1 + \sum_{i=1}^n N_{A_i}. \quad (6)$$

Equation (2) can be rewritten in the form

$$\frac{1}{z} = \mathcal{F}(\mathcal{Z}) \quad (7)$$

$$\mathcal{F}(x) = x^{-1} \left(1 + \sum_{n=1}^{\infty} \alpha_n x^n \right). \quad (8)$$

For small enough z the solution $\mathcal{Z}(z)$ is an analytic function but at some critical value $z = z_{\text{cr}}$ it is non-analytic; this is the point at which the graphical expansion (3) diverges. The susceptibility is given by

$$\chi(z) = \frac{\partial \mathcal{Z}}{\partial z} = -\frac{z^{-2}}{\mathcal{F}'(\mathcal{Z}(z))} = \frac{\mathcal{Z} z^{-1}}{1 - z \sum_{n=1}^{\infty} n \alpha_n \mathcal{Z}(z)^{n-1}} \quad (9)$$

and so the critical point is where $\mathcal{F}'(\mathcal{Z}(z_{\text{cr}})) = 0$. As $z \rightarrow z_{\text{cr}}$ the susceptibility has leading non-analytic behaviour given by

$$\chi(z) \simeq (z_{\text{cr}} - z)^{-\gamma} \quad (10)$$

where γ is the susceptibility exponent. (Throughout this letter we use the symbol $\simeq$ to denote the leading singularity as $z \rightarrow z_{\text{cr}}$.) The nature of the critical point and the value of γ depend upon how $\mathcal{F}'$ vanishes. The multi-critical BPs are obtained

by supposing that $\mathcal{F}^{(n)}(\mathcal{Z}(z_{\text{cr}})) = 0$, $n = 0, \dots, k$ but that $\mathcal{F}^{(k+1)}(\mathcal{Z}(z_{\text{cr}})) \neq 0$ so, close to z_{cr} ,

$$\chi(z) \simeq \frac{z^{-2}}{|\mathcal{Z}(z) - \mathcal{Z}(z_{\text{cr}})|^k} . \quad (11)$$

However

$$\mathcal{Z}(z) \simeq \mathcal{Z}(z_{\text{cr}}) - \text{const.} (z_{\text{cr}} - z)^{1-\gamma} \quad (12)$$

so comparing singularities we obtain

$$\begin{aligned} \gamma &= k(1 - \gamma) \\ \text{or } \gamma &= 1 - \frac{1}{k+1} . \end{aligned} \quad (13)$$

The generic case, requiring no special tuning of the α_n (except that at least one α_n ($n \geq 2$) must be non-zero), is $k = 1$, $\gamma = \frac{1}{2}$; higher values of k , which require at least k of the α_n ($n \geq 2$) to be non-zero and of varying sign to enforce the vanishing of higher derivatives, give the multi-critical BPs (MCBP from now on).

The MCBPs are not the only ones with $\gamma \neq \frac{1}{2}$. By allowing an infinite number of the α_n to be non-zero we can arrange that $\mathcal{F}'$ vanishes non-analytically [4, 5]. Consider

$$\mathcal{F}(\mathcal{Z}) = \frac{1}{\mathcal{Z}} + \mu + \frac{\mathcal{Z}}{A^2} + \lambda \left(1 - \frac{\mathcal{Z}}{A}\right)^{\beta+1} \quad (14)$$

where A , λ , μ and β are positive constants. Then

$$\mathcal{F}'(\mathcal{Z}) = -\frac{1}{\mathcal{Z}^2} + \frac{1}{A^2} - \frac{\lambda(\beta+1)}{A} \left(1 - \frac{\mathcal{Z}}{A}\right)^{\beta} . \quad (15)$$

As z increases from zero so does $\mathcal{Z}(z)$. The first zero of $\mathcal{F}'$ is at $\mathcal{Z} = A$ when $z_{\text{cr}} = (2/A + \mu)^{-1}$; provided that $1 > \beta > 0$ the non-analytic term in (15) dominates as $z \rightarrow z_{\text{cr}}$. Inserting this dominant behaviour into (11), using (12), and comparing singularities, we obtain

$$\gamma = \frac{\beta}{\beta+1} . \quad (16)$$

Thus we get models with continuously varying γ in the range 0 to $\frac{1}{2}$ as discussed in [5]. However the restriction on the range of β is easily removed; by tuning the coefficients α_n so that the analytic part of $\mathcal{F}'$ in (15) vanishes as $(\mathcal{Z} - A)^2$ the non-analytic behaviour dominates for $2 > \beta > 0$ so now we can get continuously varying γ s up to $\frac{2}{3}$. If the coefficients are tuned so that the analytic part of $\mathcal{F}'$ in (15) vanishes as $(\mathcal{Z} - A)^m$ then β can be as large as m . In this way we see that by combining the multi-critical strategy and the non-analytic form of $\mathcal{F}$ used in [4, 5] we can obtain all γ values in the range 0 to 1. If β is an integer these models just

reproduce the MCBP; when β is not an integer we will call them the ‘continuous critical branched polymers’, or CCBP.

The CCBPs actually continue to negative gamma. Suppose that $\mathcal{F}'$ is finite at z_{cr} but that $\mathcal{F}''$ diverges; then by (9) the susceptibility χ is finite at z_{cr} . However

$$\frac{\partial \chi}{\partial z} = \frac{2z^{-3}}{\mathcal{F}'(\mathcal{Z}(z))} + \frac{z^{-2}\mathcal{F}''(\mathcal{Z}(z))\chi}{\mathcal{F}'(\mathcal{Z}(z)^2)} \quad (17)$$

so the derivative of χ diverges at the critical point and γ is negative. This is easily arranged by (for example) eliminating the linear term in (14) so that

$$\mathcal{F}(\mathcal{Z}) = \frac{1}{\mathcal{Z}} + \mu + \lambda \left(1 - \frac{\mathcal{Z}}{A}\right)^{\beta+1} \quad (18)$$

with $1 > \beta > 0$ and comparison of the behaviour as $z \rightarrow z_{\text{cr}}$ then gives

$$\gamma = \frac{-\beta}{2-\beta}. \quad (19)$$

In our calculation of the spectral dimension we will need to know the behaviour of

$$\Omega_l = \frac{d^{l+1}}{d\mathcal{Z}^{l+1}} \sum_{n=1}^{\infty} \alpha_n \mathcal{Z}(z)^n = \frac{d^{l+1}}{d\mathcal{Z}^{l+1}} \mathcal{Z}\mathcal{F}(\mathcal{Z}) \quad (20)$$

close to the critical point. Using the properties of $\mathcal{F}$ discussed above we find that for $\gamma > 0$

$$\begin{aligned} \Omega_l &\simeq (\mathcal{Z}(z) - \mathcal{Z}(z_{\text{cr}}))^{\beta-l} + \frac{1}{z_{\text{cr}}} \delta_{l,0} \\ &= (z_{\text{cr}} - z)^{\gamma-l(1-\gamma)} + \frac{1}{z_{\text{cr}}} \delta_{l,0} \end{aligned} \quad (21)$$

unless β is an integer (ie MCBP) in which case for $l \geq \beta$ we find that Ω_l is finite at z_{cr} . If $\gamma < 0$ then $\mathcal{F}'$ is finite and it follows from (9) that

$$1 - z_{\text{cr}}\Omega_0(z_{\text{cr}}) = C \quad (22)$$

where C is finite.

Our calculation of the spectral dimension follows the same method as in [2] but generalized to take account of the presence of vertices of varying order. The return probability generating function $P_A(y)$ on a given polymer $A \in \mathcal{B}^n$ is related to that on its n constituents $A_1, \dots, A_n$ (see fig.1). It is convenient to define for any polymer $A \in \mathcal{B}$

$$P_A(y) = \frac{1}{1-y} \frac{1}{h_A(y)} \quad (23)$$

and then we find for $A \in \mathcal{B}^n$ [2]

$$h_A(y) = \frac{1 + \sum_{i=1}^n h_{A_i}(y)}{1 + (1-y) \sum_{i=1}^n h_{A_i}(y)}. \quad (24)$$

Note that at $y = 1$ the solution is

$$h_A(1) = N_A \quad (25)$$

for all polymers $A \in \mathcal{B}$. The spectral dimension is found by considering the quantity

$$\tilde{\mathcal{Q}}(z, y) = -\frac{d}{dy}(1-y) \sum_{A \in \mathcal{B}} z^{N_A} w_A P_A(y) = \frac{d}{dy} \sum_{A \in \mathcal{B}} \frac{z^{N_A} w_A}{h_A(y)}. \quad (26)$$

At $y = 1$ the n th derivative with respect to y of $\tilde{\mathcal{Q}}(z, y)$ behaves as

$$\tilde{\mathcal{Q}}_n(z) = \left. \frac{\partial^n}{\partial y^n} \tilde{\mathcal{Q}}(z, y) \right|_{y=1} \simeq (z_{\text{cr}} - z)^{\beta - n\Delta} \quad (27)$$

where the exponents β and Δ are related to the spectral dimension by

$$\beta = 1 - \gamma + \Delta(d_s/2 - 1). \quad (28)$$

The behaviour (27) can be established by considering the quantities

$$H^{(n_1, n_2, \dots, n_p)} = \sum_{A \in \mathcal{B}} z^{N_A} w_A \prod_{i=1}^p \left. \frac{d^n}{dy^n} h_A(y) \right|_{y=1} \quad (29)$$

which have leading singular behaviour as $z \rightarrow z_{\text{cr}}$ given by [2]

$$H^{(n_1, n_2, \dots, n_p)} \simeq (z_{\text{cr}} - z)^{a - bp - c \sum_{i=1}^p n_i} \quad (30)$$

where we will determine the constants a, b, c .

First we will consider the case when $\gamma > 0$. Let us compute

$$\begin{aligned} H^{(1)} &= \sum_{A \in \mathcal{B}} z^{N_A} w_A h'_A(1) \\ &= \sum_n \sum_{A \in \mathcal{B}^n} z^{N_A} w_A h'_A(1). \end{aligned} \quad (31)$$

Now we use (24) to relate h'_A to the corresponding quantity for the constituents of A and (4) to relate the weights and obtain

$$\begin{aligned} H^{(1)} &= \sum_n \alpha_n \sum_{A \in \mathcal{B}^n} \sum_{A_1 \cup A_2 \cup \dots \cup A_n = A} z^{1+N_{A_1}+\dots+N_{A_n}} w_{A_1} \dots w_{A_n} \\ &\quad \left\{ \sum_{i=1}^n h'_{A_i}(1) + \sum_{i=1}^n h_{A_i}(1) \left(1 + \sum_{j=1}^n h_{A_j}(1) \right) \right\}. \end{aligned} \quad (32)$$

Note that although $A \in \mathcal{B}^n$ in the above expression there is no such restriction on its constituent polymers, $A_i \dots A_n$ which are drawn from the entire ensemble $\mathcal{B}$. The terms on the r.h.s. of (32) which involve $h'_{A_m}(1)$, of which there are n , simply reproduce $H^{(1)}$ multiplied by powers of the partition function; by using (25) we see that the remaining pieces simply produce derivatives of the partition function so we may rearrange (32) to obtain

$$H^{(1)} \left(1 - z \sum_{n=1}^{\infty} n \alpha_n \mathcal{Z}(z)^{n-1} \right) = z^2 \frac{\partial^2}{\partial z^2} \mathcal{Z}. \quad (33)$$

The coefficient of $H^{(1)}$ is $1 - z\Omega_0$ so by (10) and (21) we obtain

$$H^{(1)} \simeq (z_{\text{cr}} - z)^{-1-2\gamma} \quad (34)$$

and also, since $H^{(1)}$ is the second derivative of $\tilde{\mathcal{Q}}_0(z)$ with respect to z , that

$$\tilde{\mathcal{Q}}_0(z) \simeq (z_{\text{cr}} - z)^{1-2\gamma} \quad (35)$$

which gives $\beta = 1 - 2\gamma$.

Next we compute

$$\begin{aligned} H^{(1,1)} &= \sum_{A \in \mathcal{B}} z^{N_A} w_A h'_A(1) h'_A(1) \\ &= \sum_n \alpha_n \sum_{A \in \mathcal{B}^n} \sum_{A_1 \cup A_2 \cup \dots \cup A_n = A} z^{1+N_{A_1} + \dots + N_{A_n}} w_{A_1} \dots w_{A_n} \left\{ \sum_{i=1}^n h'_{A_i}(1) \right\}^2 \\ &\quad + 2z^2 \frac{\partial^2}{\partial z^2} \sum_n \alpha_n \sum_{A \in \mathcal{B}^n} \sum_{A_1 \cup A_2 \cup \dots \cup A_n = A} z^{1+N_{A_1} + \dots + N_{A_n}} w_{A_1} \dots w_{A_n} \left\{ \sum_{i=1}^n h'_{A_i}(1) \right\} \\ &\quad + \left(z^2 \frac{\partial^2}{\partial z^2} \right)^2 \mathcal{Z}. \end{aligned} \quad (36)$$

Again the first sum contains pieces which reproduce $H^{(1,1)}$; the remaining terms on the r.h.s. of (36) can be expressed in terms of $H^{(1)}$ and $\mathcal{Z}$. We obtain

$$\begin{aligned} H^{(1,1)} \left(1 - z \sum_{n=1}^{\infty} n \alpha_n \mathcal{Z}(z)^{n-1} \right) &= z \sum_{n=2}^{\infty} n(n-1) \alpha_n \mathcal{Z}^{n-2} (H^{(1)})^2 \\ &\quad + 2z^2 \frac{\partial^2}{\partial z^2} z \sum_{n=1}^{\infty} n \alpha_n \mathcal{Z}(z)^{n-1} H^{(1)} \\ &\quad + \left(z^2 \frac{\partial^2}{\partial z^2} \right)^2 \mathcal{Z}. \end{aligned} \quad (37)$$

Noting that the coefficient of the $(H^{(1)})^2$ term is simply $z\Omega_1$ and that the coefficient of the $H^{(1)}$ term is $z\Omega_0$ which tends to 1 as $z \rightarrow z_{\text{cr}}$ we see that the first two terms on

the r.h.s. of (37) both vary as $(z_{\text{cr}} - z)^{-3-2\gamma}$ whilst the last term is sub-leading since it varies as $(z_{\text{cr}} - z)^{-3-\gamma}$. The coefficient of $H^{(1,1)}$ varies, as before, like $(z_{\text{cr}} - z)^\gamma$ so we find

$$H^{(1,1)} \simeq (z_{\text{cr}} - z)^{-3-3\gamma}. \quad (38)$$

We can iterate this process; $H^{(1,1,1)}$ is given by

$$\begin{aligned} H^{(1,1,1)}(1 - z\Omega_0) &= z\Omega_2 (H^{(1)})^3 + z\Omega_1 H^{(1,1)} H^{(1)} + 3z^2 \frac{\partial^2}{\partial z^2} z\Omega_0 H^{(1,1)} \\ &\quad + (\text{less singular terms}). \end{aligned} \quad (39)$$

The presence of the coefficients $\Omega_{1,2}$ in the first two terms on the r.h.s. changes their degree of divergence to $(z_{\text{cr}} - z)^{-5-3\gamma}$ which is the same as that of the third term and hence we conclude that

$$H^{(1,1,1)} \simeq (z_{\text{cr}} - z)^{-5-4\gamma}. \quad (40)$$

This process may be continued to longer strings $(1, 1, 1, \dots, 1)$ and to strings involving higher derivatives of $h_A(y)$ (obtained by successive differentiation of (24)). Following [2] we can set up a proof by induction [6] that

$$H^{(n_1, n_2, \dots, n_p)} \simeq (z_{\text{cr}} - z)^{1-\gamma-p-(1+\gamma)\sum_{i=1}^p n_i} \quad (41)$$

from which it follows, by inserting this result in (27) (see [2] for details), that $\Delta = 1 + \gamma$. Substituting this and $\beta = 1 - 2\gamma$ into (28) we find that

$$d_s = \frac{2}{1 + \gamma}. \quad (42)$$

Note that the spectral dimension also satisfies the scaling relation $d_s \Delta = 2$. As was discussed in [7] this scaling relation is expected to be true for graphs whose Laplacian has no level-dependent degeneracy in its eigenvalue spectrum; clearly the BP ensembles, most of whose graphs have no symmetries, fall into this category.

The case of $\gamma < 0$ is simpler. The factor $1 - z\Omega_0$ does not vanish as $z \rightarrow z_{\text{cr}}$ so we now find that

$$H^{(1)} \simeq (z_{\text{cr}} - z)^{-1-\gamma} \quad (43)$$

and therefore that

$$\tilde{\mathcal{Q}}_0(z) \simeq (z_{\text{cr}} - z)^{1-\gamma} \quad (44)$$

which implies $\beta = 1 - \gamma$. Using (28) we see immediately that $d_s = 2$. Of course it is necessary to check that $\Delta \neq 0$ before reaching this conclusion; in fact it is straightforward to show that $\Delta = 1$ [6] and so the scaling relation $d_s \Delta = 2$ is still satisfied.

It is interesting to compare these results with a scaling relation recently found by Ambjørn et al [8]. They showed that

$$\frac{2}{d_H} = \Delta \left(1 - \frac{d_s}{2}\right) \quad (45)$$

where d_H is the *extrinsic* Hausdorff dimension. For the k th multicritical model d_H has been calculated [3] and is known to be given by

$$d_H = \frac{2(k+1)}{k}. \quad (46)$$

Assuming that $d_s \Delta = 2$ we can then use (45) to determine that [8]

$$d_s = \frac{2}{2 - \frac{1}{k+1}} = \frac{2}{1 + \gamma} \quad (47)$$

in agreement with our calculations. The value of d_H has not been calculated explicitly for the CCBPs but we can now use the scaling relation (45) and our results to show that $\gamma d_H = 2$ if $\gamma > 0$. When $\gamma < 0$ we have $d_s = 2$ so we expect $d_H = \infty$. It is amusing that the two-dimensional quantum gravity models at $c < 1$ also have $d_H = \infty$ and hence $d_s = 2$ [8].

We acknowledge valuable conversations with Jan Ambjørn who told us of the scaling relation result (45) prior to writing the paper [8]. J.C. acknowledges a grant from PRAXIS XXI.

References

- [1] J.Ambjørn, J.Jurkiewicz and Y.Watabiki, Nucl. Phys. B454 (1995) 313.
- [2] T. Jonsson and J.F. Wheeler, “The spectral dimension of the branched polymer phase of two dimensional quantum gravity”, Oxford University Preprint OUTP-97-33P, hep-lat/9710024.
- [3] J.Ambjørn, B.Durhuus and T.Jonsson, Phys. Lett. B244 (1990) 403.
- [4] J.Ambjørn, B. Durhuus, J. Fröhlich and P.Orland, Nucl. Phys. B270 [FS16] 457.
- [5] P. Bialas and Z. Burda, Phys. Lett. B 384 (1996) 75.
- [6] J.D. Correia, D.Phil Thesis, Oxford University 1998.

- [7] T. Jonsson and J.F. Wheeler, “The spectral dimension on branched polymer ensembles”, to appear in the proceedings of NATO workshop *New Developments in Quantum Field Theory*, eds. P.H. Damgaard and J. Jurkiewicz, published by Plenum Publishing Corp., New York.
- [8] J. Ambjørn, D. Boulatov, J.L. Nielsen, J. Rolf and Y. Watabiki, “The spectral dimension of 2d quantum gravity”, Niels Bohr Institute preprint NBI-HE-97-62.